

# MRI NEURAL NETWORK SEGMENTATION IN ATHEROSCLEROSIS

---

[Jha, Abhinav Kumar](#), [Li, Ran](#), [Woodard, Pamela](#), [Zheng, Jie](#)

[Markiewicz, Gregory](#)

T-020254

## Technology Description

Researchers at Washington University in St. Louis have developed a two-stage neural network model, with CNN and BNN architecture, to segment carotid atherosclerotic plaque components based on multi-weighted MR images and measure the uncertainty of the segmentation output. This model identifies the lipid-rich necrotic core of the carotid atheroma for use in determining the plaque's vulnerability to rupture and cause ischemic stroke.

## Stage of Research

Researchers have trained the networks using high-resolution MRI ex vivo data, as well as pathology samples of the same plaque obtained from patients post-surgery.

## Publications

- Li R, Zheng J, Zayed MA... Jha AK. (2023). Carotid atherosclerotic plaque segmentation in multi-weighted MRI using a two-stage neural network: advantages of training with high-resolution imaging and histology. *Frontiers in Cardiovascular Medicine*, 10:1127653.

## Applications

- Diagnostic imaging for potential stroke risk

## Key Advantages

- Reliable and automated segmentation method

**Patents:** Pending

**Related Web Links:** Woodard [Profile](#) & [Lab](#)